




Objective:

An \_\_\_\_\_ can be formed by rotating one ray away from a fixed ray indicated by an arrow. The \_\_\_\_\_ is the **initial side** and the rotated ray is the \_\_\_\_\_. An angle whose vertex is the center of a circle is a \_\_\_\_\_, and the arc of the circle through which the terminal side moves is the \_\_\_\_\_. An angle in \_\_\_\_\_ is located in a rectangular coordinate system with the vertex at the origin and the initial side on the positive  $x$ -axis.



Greek letters used for the variable of an unknown angle:

$\alpha$  = alpha     $\beta$  = beta     $\gamma$  = gamma     $\theta$  = theta

**Degree Measure of Angles**






The measure,  $m(\alpha)$ , of an angle  $\alpha$  is the amount of rotation from the initial side to the terminal side, and is found by using any circle centered at the vertex. An angle that forms a complete circle arc is  $360^\circ$ .

The **degree measure of an angle** is the number of degrees in the intercepted arc of a circle centered at the vertex.

**Counterclockwise rotation**— \_\_\_\_\_

**Clockwise rotation**— \_\_\_\_\_

An angle in standard position is said to lie in the quadrant where its terminal side lies.



**Acute angle**—

**Obtuse angle**—

**Right angle**—

**Straight angle**—

**Quadrantal Angle**—

**Example:** Draw each angle in standard position, then name the quadrant in which the terminal side lies.

a)  $255^\circ$

b)  $-650^\circ$

c)  $1360^\circ$

How do you measure the line? \_\_\_\_\_

[https://commons.wikimedia.org/wiki/File:Circle\\_radians.gif#/media/File:Circle\\_radians.gif](https://commons.wikimedia.org/wiki/File:Circle_radians.gif#/media/File:Circle_radians.gif)

**Unit circle:** A circle of radius one that is centered at the origin.

What is the circumference of the unit circle? (Circumference of a circle:  $C = \pi d = 2\pi r$ )

What is the arc length intercepted by a  $180^\circ$  angle (1/2 of the circle)?


What is the arc length intercepted by a  $120^\circ$  angle (1/3 of the circle)?

What is the arc length intercepted by a  $30^\circ$  angle? An angle of  $225^\circ$ ? An angle of  $210^\circ$ ?



You have just calculated the radian measure of each of these angles.

The \_\_\_\_\_ of the angle  $\alpha$  in standard position is the directed length of the intercepted arc on the unit circle. \_\_\_\_\_

\_\_\_\_\_ means that the radian measure is positive if the rotation of the terminal side is counterclockwise and negative if the rotation is clockwise.



**One radian:** The angle that intercepts an arc with length equal to the radius of a circle. (On the unit circle, one radian is the angle that intercepts an arc of length one.)



Since the radius of the unit circle is the real number 1, without any dimension (ft., meters, etc.), the arc length, and hence the radian measure of an angle, is also a real number without any dimension. Thus, one radian (1 rad) is the real number 1.

### Converting Between Radians and Degrees:

Since there are  $2\pi$  radians in a circle (the circumference of the unit circle is  $2\pi$ ) and  $360^\circ$  in a circle,

$$2\pi \text{ radians} = 360^\circ, \text{ or } \pi \text{ radians} = 180^\circ.$$

**Degrees**→**Radians**: multiply by

**Radians**→**Degrees**: multiply by

### Examples:

Convert the degree measures to radians:

a)  $210^\circ$


b)  $-27.2^\circ$

Convert the radian measures to degrees:

a)  $\frac{5\pi}{3}$

b) 16.7 radians

**Rectangular Coordinate System**



**Examples:** Draw each angle in standard position and determine the quadrant in which each angle lies.

a)  $-\frac{5\pi}{4}$

b)  $\frac{10\pi}{3}$

c) 13.8

d) -2.5

The terminal side of an angle may be rotated in either a positive or negative direction to get to its final position. It can also be rotated for more than one revolution in either direction. Two angles with the same terminal side are called **coterminal angles**.

**Coterminal Angles**—Angles  $\alpha$  and  $\beta$  are coterminal if and only if there is an integer  $k$  such that  $m(\beta) = m(\alpha) + 360^\circ k$  or  $m(\beta) = m(\alpha) + 2\pi k$ . To find coterminal angles in degrees, add and subtract multiples of  $360^\circ$ . To find coterminal angles in radians, add or subtract multiples of  $2\pi$ . Make sure to find a common denominator.

**Examples:** Find a positive angle and a negative angle that are coterminal with each angle:

- a)  $23^\circ$                       b)  $-146^\circ$                       c)  $\frac{7\pi}{3}$                       d)  $-\frac{\pi}{4}$                       e) 1.4

**Examples:** Determine whether the angles in each pair are coterminal:

- a)  $-128^\circ$  and  $592^\circ$                       b)  $8^\circ$  and  $-368^\circ$                       c)  $\frac{3\pi}{2}$  and  $\frac{9\pi}{2}$

What is the circumference of the unit circle? (Circumference of a circle:  $C = \pi d = 2\pi r$ )




What is the arc length intercepted by a  $180^\circ$  angle ( $1/2$  of the circle)?

What is the arc length intercepted by a  $330^\circ$  angle?

What if the radius isn't 1? How do we find the arc length of the intercepted arc?

### Arc Length

Often, we want to find the arc length,  $s$ , on a circle of radius,  $r$ , intercepted by an angle,  $\alpha$ . We can do this by determining what fraction of the circle we are looking at, then multiplying by the circumference of the circle.



**Degrees:**


**Radians:**                      or                      **This formula only works if  $\alpha$  is in radians!**

**Examples:**

Use the arc length formula and the given information to find the arc length. If you are given radians, write the answer in terms of  $\pi$  and use the  $\pi$  button to round answers to the nearest tenth. If you are given degrees, round the answer to the nearest tenth of a degree. Show work using the formula. Remember, leave work and answer in the units it begins with.


a.  $r = 5 \text{ ft.}, \theta = 150^\circ$

b.  $r = 2.5 \text{ km}, \theta = \frac{5\pi}{3}$



Use the arc length formula and the given information to find the indicated measure. Round answers to the nearest tenth if necessary. Show work using the formula. Remember, leave work and answer in the units it begins with.

c.  $s = 3.5 \text{ ft.}, \theta = \frac{3\pi}{4} \text{ rad};$  find  $r$



d.  $r = 3 \text{ in.}, s = 20.5;$  find  $\theta$  in (leave answer in degrees)